

SIMULTANEOUS ELECTRON SPECTROSCOPY AND X-RAY SCATTERING ON MODEL CERIA CATALYSTS

Baran Eren

*Department of Chemical and Biological Physics, Weizmann Institute of Science, 234 Herzl Street, 76100
Rehovot, Israel*

Heterogeneous catalysis is a timely and critical research field in basic and applied energy sciences, due to its potential to provide solutions to global environmental issues. However, there is still a lack of a profound understanding of the molecular and structural processes at the interfaces between solids and reactant gases. A detailed understanding of the correlation between the chemistry, structure, and function in these materials requires a multimodal investigation. Over the past few years, scientists at the Advanced Light Source, the Berkeley synchrotron facility, have developed a unique setup attached to an X-ray beamline where chemically-sensitive ambient pressure X-ray photoelectron spectroscopy (APXPS) and structure-sensitive grazing incidence X-ray scattering (GIXS) experiments can be performed simultaneously. Here, we showcase that this tool can provide mechanistic insights that are unparalleled in the literature.

Our novel approach allows us to probe the changing surface and bulk chemistry, and surface and bulk structure of the model ceria catalysts in the presence of H₂ and CO₂ gases. In addition to the method itself, the electron density, surface chemistry, and roughness trends observed in ceria during the reaction will be discussed in this talk. Access to such a variety of data from working catalysts in a single experiment can have far-reaching implications, because changes in surface roughness, ability to store hydrogen in the bulk in various forms, and the chemical state of the surface, which all depend on the reactive environment, can directly affect the catalyst performance.